Carnegie Veterinary Clinic & Carnegie Cat Cottage



66 Rosstown Road Carnegie, Victoria 3163 P: 9563 5422 F: 9563 6944



Email: cvc66000@commander360.com Website: www.carnegievets.com.au & www.carnegiecatcottage.com.au

Our Vets:

Dr. Ketan Jivanjee

BSc (La Trobe), BVSc (Pretoria), MRCVS (UK) Special Interests: Small Animal Surgery, Small Animal Internal Medicine, Ophthalmology, Theriogenology, Pharmacology, Unusual & Exotic Pet Medicine & Surgery, Veterinary Practice Management.

Dr. Kelly Wilkinson

BVSc (Hons) (Melbourne)

Special interests: Small Animal Medicine, Feline Medicine, Animal Behaviour, Unusual & Exotic Pet Medicine & Surgery.

You may have already met our nurses; Natasha, Shonna, Amy and Melissa. They have pets of their own and are well versed on all clinic, hospital and cat boarding services. Please do not hesitate to ask them any questions. They will assist you as much as possible to ensure a well informed and worthwhile visit to the clinic. Our staff will eagerly assist you with your appointments, food supplies and any other questions you may have about the care and welfare of your pet. Passion, Pride and Perseverance in Service and Care is what we do.

Natasha Whitehead Cert IV (Vet. Nursing) Areas of interest: Surgical nursing, Practice Management, Puppy Preschool, Nutrition & Behaviour, Counselling & Client Relations

Shonna Camilleri Cert IV (Vet. Nursing) Areas of interest: Surgical Nursing, Medicial Nursing, Nutrition & Behaviour, Counselling & Client Relations

Sajani De Silva *Studying for Cert IV (Vet. Nursing)* Areas of interest: Surgical Nursing, Medicial Nursing, Nutrition, Counselling & Client Relations

Melissa Collins

Areas of interest: Nursing, Client Relations, Cattery Management

Carnegie Veterinary Clinic Hours:

Monday to Friday: 8am – 7pm Saturday 9am – 1:30pm

Carnegie Cat Cottage – Hours of Admission and Discharge:

Monday to Friday: 10am – 1pm Saturday 9am 1:30pm

In the unfortunate event that you would require veterinary assistance after hours please contact:

Pet Emergency Centre, 1103 Dandenong Road, Malvern East, Victoria 3145

Phone: 9569 3677 Melways Reference: 68 J3 **Southern Animal Emergency Centre**, 248 Wickham Road, Highett, Victoria 3190 Phone:9532 5261 Melways Reference: 77 H8

Feline Immunodeficiency Virus infection needs special care

Feline Immunodeficiency Virus (FIV) is similar to the Human Immunodeficiency Virus (HIV) in humans. Like HIV it may or may not proceed to cause clinical illness in the cats it infects. FIV is transmitted primarily via saliva and the most common causes are fighting and biting. FIV infection can predispose cats to numerous problems including infections (fungal, bacterial and parasitic), bone marrow disease, oral disease, cancer and eye problems. A vaccine is now available for kittens or older cats, but your cat should be confirmed clear of infection before its use.

Diagnosis usually relies on the detection of *antibodies* to the virus in the cat's blood sample. It is rare to get a "false-positive" result, however nursing kittens of FIV-positive mums may test positive for some time. They may not be truly infected with the virus but have antibodies from the mother's milk.

If your cat has been in a fight, it is wise to wait at least 60 days before testing for FIV in order to minimise the risk of a falsely negative result. While a positive test result usually indicates infection, a negative result could indicate that either the cat is not infected, or it has been exposed but is not yet producing antibodies. A negative result may also occur if the cat has been immunosuppressed by the virus and cannot produce antibodies, or if the test itself is faulty.

Management of cats with FIV requires some adjustments to their lifestyle. These cats should be kept indoors to minimise spread to other cats AND to decrease exposure to disease. Ideally, infected cats will have a full clinical examination twice a year. Your vet will usually spend time examining the mouth, lymph nodes and eyes, as these are all areas that can be affected

by the disease. Your cat will often be weighed as weight loss can be an early warning sign of clinical disease. Some vets will also recommend an annual blood profile, and occasionally urine or faeces tests. Generally, vaccination programs for FIV-infected cats should be similar to those for uninfected cats, however vets will often use a 'killed' vaccine where available.

When a cat is diagnosed with FIV-related disease, treatment decisions (including euthanasia) should not be made solely on the grounds of the cat having FIV. Similarly, diseases in cats infected with FIV may not necessarily be a result of that infection, but may have occurred independently. Overall, many FIV infected cats, will live long and happy lives regardless of their infection status.



Enter our competition and WIN! Details above

- Pregnancy diagnosis in dogs and cats
- Precocious guinea pigs
- Feline hyperthyroidism



False pregnancy in the bitch

It is possible for any bitch that has not been spayed to exhibit symptoms of pregnancy without actually being pregnant. Some bitches can completely fool their owners with convincing displays of maternal behaviour. Just like a real pregnancy, false-pregnant bitches gain weight, produce milk and make nests. They may be listless, off their food and will often "mother" toys.



It is unclear as to why some bitches experience false pregnancies, but we do know that any bitch that ovulates, whether pregnant or not, will experience the same blood levels of a hormone called progesterone. Around two months after ovulation, the progesterone levels drop and the "milk" hormone prolactin starts to increase in concentration. It is prolactin that is responsible for milk production and maternal behaviour.

False pregnancies generally run a course of 2-3 weeks and treatment is not usually required. In severe cases, where large amounts of milk are being produced, a reduction in daily food intake may help the milk supply to drop. The bitch must not

be allowed to lick at her nipples as any stimulation of the mammary glands promotes milk production.

False pregnancies are a normal phenomenon and are not associated with fertility problems. They frequently recur but can be prevented with sterilisation. If the bitch is intended for breeding then she should be bred from at her next season and then sterilised once her breeding days are over.

Desensitise senstive areas

Many animals have particular areas of their body that they consider private and would prefer to keep that way. Unfortunately as Veterinary Surgeons we are often forced to invade this privacy and go into those forbidden zones. Taking a rectal temperature, clipping nails, examining down ears and checking teeth can cross the boundary for some animals. To help visits to the vet be smoother and less traumatic for your pet we would recommend a gentle desensitisation program. Handle your pet's sensitive areas regularly – ideally daily. Pick up their feet and gently finger between the toes, lift up ear flaps, handle around the lips and gums and raise the tail. This can be disguised as play and rewarded with food treats and verbal praise.



Precocious guinea pigs

With most animals, the general consensus is to wait until they are fully grown and mature before allowing them to breed. opposite is true with guinea pigs. If you do wish to breed your female guinea pig, it is important to do so BEFORE she is fully mature. This is because guinea pigs produce relatively enormous offspring. In order to deliver such a large baby, the sow's pelvic canal must enlarge by an extra 1.5 to 3 cm. A young guinea pig's pelvic bones will separate easily and the resulting gap will fill with fibro-cartilage (gristle).



By the time a sow is 9 months old, her pelvis will be fully formed. The small joints or suture lines in the pelvis, where separation was possible, will now be fused with solid bone. The resulting birth canal is usually too narrow to allow a natural delivery and the sow gets into birthing difficulties.

The only treatment in this situation is a caesarean section under general anaesthetic - a risky procedure in an animal predisposed to anaesthetic complications. To prevent this scenario, breed from your sow prior to 7 months of age and you will be able to enjoy her wonderfully precocious offspring with minimal stress.



Feline hyperthyroidism

Feline hyperthyroidism is one of the most common diseases affecting middle aged to older cats. Affected cats have over-active thyroid glands that produce too much thyroid hormone. This is usually caused by over-functioning nodules within the thyroid gland. Occasionally, in 1-2% of cases, it is secondary to a cancer of the thyroid. Since thyroid hormone affects multiple organs and helps to control an animal's metabolic rate, there can be many different presenting symptoms.

Hyperthyroid cats usually have an increased appetite and thirst. However, despite eating more, they lose weight and condition. They often exhibit changes in behaviour and may become more cranky and agitated. Owners can be woken up at night to a loudly vocalising feline demanding a snack RIGHT NOW! Many hyperthyroid cats have intermittent vomiting and diarrhoea. As thyroid hormone also affects the heart, an increase in thyroid hormone levels will increase the heart rate and blood pressure. Left unchecked, hyperthyroidism can eventually result in heart and/or kidney failure.

Hyperthyroidism is diagnosed by blood tests. There are several treatment options that your vet will discuss with you.

Affected animals may be managed on oral medication. Sometimes they can have the affected thyroid surgically removed or undergo treatment with radioiodine. This involves the injection of a radioisotope of iodine that concentrates in and destroys abnormal thyroid tissue, while leaving normal thyroid tissue unharmed. Although this is an excellent treatment option, it is only available in a limited number of specialised facilities.

Treatment of hyperthyroidism may make underlying kidney problems worse. Therefore, your vet will need to monitor your cat's progress and occasionally may even elect not to treat the condition.



Competition for our readers! WIN a \$50 shopping voucher

1) Just answer the follow	ving question: Whic	h article did you lik	e best in this issue of
Your Vet?			

- Your Vet?
- 2) Tick the box next to your preferred voucher: ☐ K Mart Australia Ltd ☐ Coles Myer Ltd ☐ Target Australia Ltd ☐ JB Hi Fi
- 3) What is your email address? *
 - * By providing us with your email address, you agree to receive future marketing material.
- 4) Post with your name and address on the back of the envelope to: Your Vet Voucher Competition, Unit 5/1 Almondbury Road, Mt Lawley WA 6050 The winner of the competition in *Your Chiropractor's* November edition was Ms S. Coates of NSW.

THAT'S IT! GOOD LUCK!

Must be received by latest post marked 02/05/2012 to be in the running



Stimulating birds in captivity

Birds are intelligent creatures that need visual stimulation and exercise. Unfortunately, due to our busy modern lifestyle, our pet birds are often left on their own while we are at work.



Encouraging foraging behaviour helps prevent obesity and boredom. In the wild, birds must constantly forage for food. This natural behaviour provides a varied diet and exercise. In captivity, a food bowl is hardly a workout and the bird can quickly gorge until full and then has nothing to do to fill his day.

Provide a stimulating and varied diet with lots of fresh vegetables, some fruit and a complete pelleted ration. Try things such as corncobs, capsicum strips, carrot tops and wheat grass that can be hidden inside things or strung on a leather strip. Place a vase in or next to the cage and fill it with small branches and natural vegetation.

Toys can also help to create a stimulating environment for your pet bird. Provide a variety of toys and rotate them. Include 'destructible' toys, such as wooden clothes pegs and leather strips that can be chewed. You can also hide food inside toys to encourage play.

Interaction with toys can provide hours of fun, and the activity provided by toys is essential for good physical and emotional wellbeing in a captive environment. However, care should be taken to ensure that your pet bird is not swallowing pieces of indigestible toys.



Tell tails

The tail is a great mood barometer and can be the bringer of good news and bad from our pets. The perky "tail up" position is used by happy cats in combination with a bouncy trot when they greet their owners. A fluffed-up cat's tail held aloft over raised hackles indicates fear and anger. It gives the impression of a much larger animal and is used to intimidate an enemy. Prior to pouncing, a cat will hold its tail rigid and just let the tail tip twitch.

Dominant and aggressive dogs will approach a potential conflict situation with an upright tail that may be wagged slowly. This is a totally

different wag to the exuberant wag of a happy dog. Unfortunately some dogs' relentless happiness can lead to a medical condition commonly known as "happy tail". This occurs when repetitive smashing against hard surfaces chronically damages the wagging tail tip. Some of these animals end up requiring amputation of the damaged tail tip. Shy and submissive dogs use a low quick wag as they crawl along whilst really frightened dogs stop wagging all together and tuck their tail firmly between their legs.

Who needs language when you have a tail!

Cat Quiz Word Sleuth

Q	T	В	U	R	М	Е	S	Е	Ν
I	E	T	А	R	Ο	K	Ο	Α	Ν
L	S	S	W	L	Н	Y	I	М	Ν
L	Р	М	E	U	I	Ν	K	Α	R
О	Н	Α	Т	Ν	I	Ν	I	F	J
D	Y	Z	Т	S	I	S	Е	W	U
G	Ν	X	S	K	R	K	Α	S	Ο
А	X	Y	I	Е	S	Е	Ν	Α	Е
R	В	G	Р	X	Ο	Z	Е	О	В
A	Е	S	Е	М	A	I	S	W	T

CLUES

Please find the following cat breeds in the word sleuth

1. This lithe and sleek cat is one of the oldest short-haired breeds and looks similar to drawings of Ancient Egyptian cats. 2. These long-haired, active and highly talkative cats are part of the Oriental family and were given their name because their grace and elegance reminded breeders of the native dancers of Bali. 3. This is the only natural breed of brown cat. Powerfully built with a rounded chest, they are easily trained and enjoy performing tricks. 4. This cat is the only recognised hairless breed of cat in Australia. It is also the only cat that has sweat glands and therefore does not need to pant to keep cool similar to other cats and dogs. 5. This cat breed is very slender with blue almond-shaped eyes. They are often very vocal. 6. This cat originated in the USA. It is known for its gentle and docile nature and its tendency to relax and become floppy when picked up. 7. One of the most popular breeds in Australia, this cat has a long fluffy coat and a slightly squashedin facial appearance. 8. This cat breed usually has no tail. Its hind legs are also thicker and longer than the ordinary cat. 9. One of the oldest breeds of cat in the world, this silver-blue cat originated in Thailand and is considered to bring good fortune. 10. This cat breed was developed by crossing a Siamese with a Burmese.

Dr. Ketan Jivanjee

from the University of Pretoria, Onderstepoort, South Africa. Prior to that, he completed Bachelor of Science with majors in Biochemistry and Microbiology from La Trobe University, Bundoora. Ketan is a Member of the Royal College of Veterinary Surgeons, United Kingdom. In 2003, a passion for surgery led him to complete a one year post graduate course in Small Animal Surgery conducted by the University of Sydney. Ketan has focused on small animal practice since graduation. He spent three years in a

Ketan graduated Bachelor of Veterinary Science

Ketan has focused on small animal practice since graduation. He spent three years in a busy small animal hospital in Albury/Wodonga honing his medical and surgical skills. His time there was both fruitful and rewarding allowing him to provide an in-depth medical and surgical service here at Carnegie Veterinary Clinic. He enjoys the rigours, challenges and diversity of a multi-faceted general practice. Out of the clinic, Ketan partakes in indoor soccer competitions.

Dr. Kelly Wilkinson

Kelly graduated Bachelor of Veterinary Science with honours from The University of Melbourne. Kelly is passionate about feline medicine. In 2002, Kelly completed a one year post graduate course in Feline Medicine conducted by the University of Sydney. Over the years, Kelly has maintained a strong focus on continuing veterinary education. She is also a member of the European Society of Feline Medicine.

Kellie's interests away from the Clinic are her two beautiful daughters – Natalie and Stephanie.

We are passionate about continuing veterinary education and ongoing learning. All members of the Carnegie Veterinary Clinic Team partake in continuing professional development to further enhance their skills and ultimately provide better service to clients and better outcomes to patients. Striving for excellence in service and care is our priority.

To provide adequate service and facilities for your needs it is necessary to charge realistic fees for treating your pet. Prompt payment of our account will help us keep the fees down. Please note that we require complete payment at the time of each consultation visit and prior to discharge from the clinic for companions that underwent in-hospital procedures. An estimate of fees can be provided for all diagnostics and treatment. We accept cash, eftpos, mastercard and visa.

The Clinic itself does not offer an account facility. If you prefer to spread the cost of expensive and extensive medical or surgical treatments, payment plans are available through our GE Money CareCredit facility. Please ask Reception for more information.

We strongly recommend Pet Insurance. Unfortunately, a Medicare type system is not in place for our pets, so the real cost of providing treatment and care must be borne by the owner. If you would like to know more about pet insurance or Petplan specifically please feel free to pick up brochures which are readily available from reception.

Nobody likes waiting. We do our best to run on time. However, occasionally there are complex cases and emergencies, so we can run behind. We appreciate and thank you for your patience.

Answers can be found on the bottom of the inside left page.